



# Charm weak decays at BESIII and STCF

**Xiao-Rui Lyu**  
([xiaorui@ucas.ac.cn](mailto:xiaorui@ucas.ac.cn))

University of Chinese Academy Sciences

(On behalf of the BESIII collaboration and the STCF working group)



# BEPCII and STCF in China

## BEPCII

- ❑ Peak luminosity  $0.6-1 \times 10^{33} \text{ cm}^{-2}\text{s}^{-1}$  at **3.773 GeV**
- ❑ Energy range  $E_{\text{cm}} = 2 - 4.6 \text{ GeV}$
- ❑ No Polarization

## Designed STCF

- ❑ Peak luminosity  $0.5-1 \times 10^{35} \text{ cm}^{-2}\text{s}^{-1}$  at **4 GeV**
- ❑ Energy range  $E_{\text{cm}} = 2-7 \text{ GeV}$
- ❑ **Potential** to increase luminosity and realize beam polarization




- $e^+$ , a converter, a linac and a damping ring, 0.5GeV
- $e^-$ , a polarized  $e^-$  source, accelerated to 0.5GeV

**$1 \text{ ab}^{-1}$  data expected per year**

# BESIII Physics

Chinese Physics C Vol. 44, No. 4 (2020)




## Future Physics Programme of BESIII\*

**Abstract:** There has recently been a dramatic renewal of interest in hadron spectroscopy and charm physics. This renaissance has been driven in part by the discovery of a plethora of charmonium-like  $XYZ$  states at BESIII and  $B$  factories, and the observation of an intriguing proton-antiproton threshold enhancement and the possibly related  $\Lambda(1835)$  meson state at BESIII, as well as the threshold measurements of charm mesons and charm baryons. We present a detailed survey of the important topics in tau-charm physics and hadron physics that can be further explored at BESIII during the remaining operation period of BEPCII. This survey will help in the optimization of the data-taking plan over the coming years, and provides physics motivation for the possible upgrade of BEPCII to higher luminosity.

**DOI:** 10.1088/1674-1137/44/4/040001

Received 25 December 2019, Published online 26 March 2020

\* Supported in part by National Key Basic Research Program of China (2015CB856700); National Natural Science Foundation of China (NSFC) (11335008, 11425524, 11625523, 11655010, 11735014, 11822506, 11935018); the Chinese Academy of Sciences (CAS) Large-Scale Scientific Facility Program; the CAS Center for Excellence in Particle Physics (CCEPP); Joint Large-Scale Scientific Facility Funds of the NSFC and CAS (U1532257, U1532258, U1732263); CAS Key Research Program of Frontier Science (QYZDJ-SSW-SLH003, QYZDJ-SSW-SLH040); 100 Talents Program of CAS; CAS PIFI; the Thousand Talents Program of China; IN-PAC and Shanghai Key Laboratory for Particle Physics and Cosmology; German Research Foundation DFG under Contracts Nos. Collaborative Research Center CRC 1044, FOR 2359; Istituto Nazionale di Fisica Nucleare, Italy; Koninklijke Nederlandse Akademie van Wetenschappen (KNAW) (530-ACDP03); Ministry of Development of Turkey (DPT2006K-120470); National Science and Technology fund; The Knut and Alice Wallenberg Foundation (Sweden) (2016.0157); The Swedish Research Council; U. S. Department of Energy (DE-FG02-05ER41374, DESC-0010118, DE-SC-0012069); University of Groningen (RuG) and the Helmholtzzentrum fuer Schwerionenforschung GmbH (GSI), Darmstadt; the Russian Ministry of Science and Higher Education (14.W03.31.0026)

 Content from this work may be used under the terms of the Creative Commons Attribution 3.0 license. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Article funded by SCOAP<sup>3</sup> and published under licence by Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd



Int. J. Mod. Phys. A 24, S1-794 (2009)  
[arXiv:0809.1869 [hep-ex]].

Chin. Phys. C 44, 040001 (2020)  
doi:10.1088/1674-1137/44/4/040001  
[arXiv:1912.05983 [hep-ex]].

# Facilities for Charm Study

- **LHCb**: huge x-sec, boost,  $9 \text{ fb}^{-1}$  now (x40 current B-factories)
- **B-factories** (Belle(-II), BaBar): more kinematic constrains, clean environment,  $\sim 100\%$  trigger efficiency
- **$\tau$ -charm factory** : Low backgrounds and high efficiency; missing technique; Quantum correlations and CP-tagging are unique;
  - **BESIII**:  $20 \text{ fb}^{-1}$  at 3.77 GeV;  $6 \text{ fb}^{-1}$  at 4.18 GeV;  $15 \text{ fb}^{-1}$  @ 4.6-4.9 GeV
  - **STCF** :  $4 \times 10^9$  pairs of  $D^{\pm,0}$  and  $10^8$   $D_s$  pairs per year
  - **Highlighted Physics programs**
    - Precise measurement of (semi-)leptonic decay ( $f_D$ ,  $f_{D_s}$ , CKM matrix...)
    - $D^0 - \bar{D}^0$  mixing, CPV
    - Rear decay (FCNC, LFV, LNV....)
    - Excite charm meson states  $D_J$ ,  $D_{sJ}$  (mass, width,  $J^{PC}$ , decay modes)
    - Charmed baryons ( $J^{PC}$ , Decay modes, absolute BF)
    - Light meson and hyperon spectroscopy studied in charmed hadron decays

# Charmed meson decays



# Precision measurement of CKM elements



**CKM matrix** elements are **fundamental SM parameters** that describe the mixing of quark fields due to weak interaction.

- A precise test of EW theory
- New physics beyond SM?

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

Three generations of quark?

Expected precision < 2% at BESIII

Unitary matrix?

BESIII + B factories +  
LQCD


BESIII + B factories +  
LHCb + LQCD

A **direct measurement** of  $V_{cd(s)}$  is one of the most **important task** in charm physics

# $D_{(s)}$ (Semi-)Leptonic decay


## Purely Leptonic:

$$\Gamma(D_{(s)}^+ \rightarrow \ell^+ \nu_\ell) = \frac{G_F^2 f_{D_{(s)}^+}^2}{8\pi} |V_{cd(s)}|^2 m_\ell^2 m_{D_{(s)}^+}^2 \left(1 - \frac{m_\ell^2}{m_{D_{(s)}^+}^2}\right)^2$$



## Semi-Leptonic:

$$\frac{d\Gamma}{dq^2} = \frac{G_F^2}{24\pi^3} |V_{cs(d)}|^2 p_{K(\pi)}^3 |f_+^{K(\pi)}(q^2)|^2$$



**Directly measurement :**  $|V_{cd(s)}| \times f_{D(s)}$  or  $|V_{cd(s)}| \times FF$

- ❑ Input  $f_{D(s)}$  or  $f^{K(\pi)}(0)$  from LQCD  $\Rightarrow |V_{cd(s)}|$
- ❑ Input  $|V_{cd(s)}|$  from a global fit  $\Rightarrow f_{D(s)}$  or  $f^{K(\pi)}(0)$
- ❑ Validate LQCD calculation of Input  $f_{B(s)}$  and provide constrain of CKM-unitarity

# $D_{(s)}$ Leptonic decay

	BESIII	BESIII
Luminosity	2.9 fb <sup>-1</sup> at 3.773 GeV	20 fb <sup>-1</sup> at 3.773 GeV
$\mathcal{B}(D^+ \rightarrow \mu^+ \nu_\mu)$	5.1% <sub>stat.</sub> 1.6% <sub>syst.</sub> [4]	1.9% <sub>stat.</sub> 1.3% <sub>syst.</sub>
$f_{D^+}$ (MeV)	2.6% <sub>stat.</sub> 0.9% <sub>syst.</sub> [4]	1.0% <sub>stat.</sub> 0.8% <sub>syst.</sub>
$ V_{cd} $	2.6% <sub>stat.</sub> 1.0% <sub>syst.</sub> [4]	1.0% <sub>stat.</sub> 0.8% <sub>syst.</sub>
$\mathcal{B}(D^+ \rightarrow \tau^+ \nu_\tau)$	20% <sub>stat.</sub> 10% <sub>syst.</sub>	8% <sub>stat.</sub> 5% <sub>syst.</sub>
$\frac{\mathcal{B}(D^+ \rightarrow \tau^+ \nu_\tau)}{\mathcal{B}(D^+ \rightarrow \mu^+ \nu_\mu)}$	21% <sub>stat.</sub> 10% <sub>syst.</sub>	8% <sub>stat.</sub> 5% <sub>syst.</sub>
Luminosity	3.2 fb <sup>-1</sup> at 4.178 GeV	6 fb <sup>-1</sup> at 4.178 GeV
$\mathcal{B}(D_s^+ \rightarrow \mu^+ \nu_\mu)$	2.8% <sub>stat.</sub> 2.7% <sub>syst.</sub> [5]	2.1% <sub>stat.</sub> 2.2% <sub>syst.</sub>
$f_{D_s^+}$ (MeV)	1.5% <sub>stat.</sub> 1.6% <sub>syst.</sub> [5]	1.0% <sub>stat.</sub> 1.2% <sub>syst.</sub>
$ V_{cs} $	1.5% <sub>stat.</sub> 1.6% <sub>syst.</sub> [5]	1.0% <sub>stat.</sub> 1.2% <sub>syst.</sub>
$f_{D_s^+}/f_{D^+}$	3.0% <sub>stat.</sub> 1.5% <sub>syst.</sub> [5]	1.4% <sub>stat.</sub> 1.4% <sub>syst.</sub>
$\mathcal{B}(D_s^+ \rightarrow \tau^+ \nu_\tau)$	2.2% <sub>stat.</sub> 2.6% <sub>syst.</sub>	1.6% <sub>stat.</sub> 2.4% <sub>syst.</sub>
$f_{D_s^+}$ (MeV)	1.1% <sub>stat.</sub> 1.5% <sub>syst.</sub>	0.9% <sub>stat.</sub> 1.4% <sub>syst.</sub>
$ V_{cs} $	1.1% <sub>stat.</sub> 1.5% <sub>syst.</sub>	0.9% <sub>stat.</sub> 1.4% <sub>syst.</sub>
$\overline{f}_{D_s^+}^{\mu\&\tau}$ (MeV)	0.9% <sub>stat.</sub> 1.0% <sub>syst.</sub>	0.6% <sub>stat.</sub> 0.9% <sub>syst.</sub>
$ \overline{V}_{cs}^{\mu\&\tau} $	0.9% <sub>stat.</sub> 1.0% <sub>syst.</sub>	0.6% <sub>stat.</sub> 0.9% <sub>syst.</sub>
$\frac{\mathcal{B}(D_s^+ \rightarrow \tau^+ \nu_\tau)}{\mathcal{B}(D_s^+ \rightarrow \mu^+ \nu_\mu)}$	3.6% <sub>stat.</sub> 3.0% <sub>syst.</sub>	2.6% <sub>stat.</sub> 2.8% <sub>syst.</sub>

\* assuming Belle II improved systematics by a factor 2

STCF	Belle II
1 ab <sup>-1</sup> at 3.773 GeV	50 ab <sup>-1</sup> at $\Upsilon(nS)$
0.28% <sub>stat</sub>	—
0.15% <sub>stat</sub>	—
0.15% <sub>stat</sub>	—
0.41% <sub>stat</sub>	—
0.50% <sub>stat</sub>	—
1 ab <sup>-1</sup> at 4.009 GeV	50 ab <sup>-1</sup> at $\Upsilon(nS)$
0.30% <sub>stat</sub>	0.8% <sub>stat</sub> 1.8% <sub>syst</sub>
0.15% <sub>stat</sub>	—
0.15% <sub>stat</sub>	—
0.21% <sub>stat</sub>	—
0.24% <sub>stat</sub>	0.6% <sub>stat</sub> 2.7% <sub>syst</sub>
0.11% <sub>stat</sub>	—
0.11% <sub>stat</sub>	—
0.09% <sub>stat</sub>	0.3% <sub>stat</sub> 1.0% <sub>syst</sub>
0.09% <sub>stat</sub>	—
0.38% <sub>stat</sub>	0.9% <sub>stat</sub> 3.2% <sub>syst</sub>

LQCD : 0.2%  
(0.1% expected)

LQCD : 0.2%  
(0.1% expected)

LQCD : 0.2%  
(0.1% expected)

Stat. uncertainty is closed to theory precision  
Sys. is challenging





# Prospects on the semi-leptonic decays

## LFU test at STCF

Luminosity	BESIII 2.9 fb <sup>-1</sup> @3.773 GeV	BESIII 20 fb <sup>-1</sup> @3.773 GeV	Belle 0.28 ab <sup>-1</sup>	Belle II 50 ab <sup>-1</sup>
$D^0 \rightarrow K^- e^+ \nu_e$	0.4% <sub>stat.</sub> 0.5% <sub>syst.</sub>	0.2% <sub>stat.</sub> 0.4% <sub>syst.</sub>	1.0% <sub>stat.</sub> 3.2% <sub>syst.</sub> *	0.1% <sub>stat.</sub> 1.6% <sub>syst.</sub> *
$D^0 \rightarrow K^- \mu^+ \nu_\mu$	0.5% <sub>stat.</sub> 0.4% <sub>syst.</sub>	0.2% <sub>stat.</sub> 0.4% <sub>syst.</sub>		
$D^0 \rightarrow \pi^- e^+ \nu_e$	1.3% <sub>stat.</sub> 0.7% <sub>syst.</sub>	0.5% <sub>stat.</sub> 0.4% <sub>syst.</sub>	3.2% <sub>stat.</sub> 4.8% <sub>syst.</sub> *	0.2% <sub>stat.</sub> 2.4% <sub>syst.</sub> *
$D^0 \rightarrow \pi^- \mu^+ \nu_\mu$	NA	0.8% <sub>stat.</sub> 0.8% <sub>syst.</sub>		
$D^0 \rightarrow K^{*-} e^+ \nu_e$				
$r_V$	5.0% <sub>stat.</sub> 2.0% <sub>syst.</sub>	2.0% <sub>stat.</sub> 2.0% <sub>syst.</sub>	—	—
$r_A$	10.% <sub>stat.</sub> 2.0% <sub>syst.</sub>	4.0% <sub>stat.</sub> 2.0% <sub>syst.</sub>	—	—
$D^0 \rightarrow a_0^-(980) e^+ \nu_e$	NA	10.% <sub>stat.</sub> 5.0% <sub>syst.</sub>	—	—
$D^0 \rightarrow K_1^-(1270) e^+ \nu_e$	NA	10.% <sub>stat.</sub> 5.0% <sub>syst.</sub>	—	—
$D^+ \rightarrow K^0 e^+ \nu_e$	0.6% <sub>stat.</sub> 1.7% <sub>syst.</sub>	0.2% <sub>stat.</sub> 1.0% <sub>syst.</sub>	—	—
$D^+ \rightarrow K_L^0 e^+ \nu_e$	0.9% <sub>stat.</sub> 1.6% <sub>syst.</sub>	0.4% <sub>stat.</sub> 1.0% <sub>syst.</sub>	—	—
$D^+ \rightarrow K^0 \mu^+ \nu_\mu$	NA	0.3% <sub>stat.</sub> 1.0% <sub>syst.</sub>	—	—
$D^+ \rightarrow \bar{K}^{*0} e^+ \nu_e$				
$A_1(0)$	1.7% <sub>stat.</sub> 2.0% <sub>syst.</sub>	0.7% <sub>stat.</sub> 1.0% <sub>syst.</sub>	—	—
$r_V$	4.0% <sub>stat.</sub> 0.5% <sub>syst.</sub>	1.6% <sub>stat.</sub> 0.5% <sub>syst.</sub>	—	—
$r_A$	5.0% <sub>stat.</sub> 1.0% <sub>syst.</sub>	2.0% <sub>stat.</sub> 1.0% <sub>syst.</sub>	—	—
$D^+ \rightarrow \pi^0 e^+ \nu_e$	1.9% <sub>stat.</sub> 0.5% <sub>syst.</sub>	0.7% <sub>stat.</sub> 0.5% <sub>syst.</sub>	—	—
$D^+ \rightarrow \pi^0 \mu^+ \nu_\mu$	NA	1.0% <sub>stat.</sub> 1.0% <sub>syst.</sub>	—	—
$D^+ \rightarrow \eta e^+ \nu_e$	4.5% <sub>stat.</sub> 2.0% <sub>syst.</sub>	2.0% <sub>stat.</sub> 2.0% <sub>syst.</sub>	—	—
$D^+ \rightarrow \eta' e^+ \nu_e$	NA	10.% <sub>stat.</sub> 5.0% <sub>syst.</sub>	—	—
$D^+ \rightarrow \omega e^+ \nu_e$				
$r_V$	7.2% <sub>stat.</sub> 4.8% <sub>syst.</sub>	3.0% <sub>stat.</sub> 2.0% <sub>syst.</sub>	—	—
$r_A$	14% <sub>stat.</sub> 5.0% <sub>syst.</sub>	3.0% <sub>stat.</sub> 2.0% <sub>syst.</sub>	—	—
$D^+ \rightarrow a_0^0(980) e^+ \nu_e$	NA	10.% <sub>stat.</sub> 5.0% <sub>syst.</sub>		
$D^+ \rightarrow \bar{K}_1^0(1270) e^+ \nu_e$	NA	10.% <sub>stat.</sub> 5.0% <sub>syst.</sub>		
$D^{0(+)} \rightarrow \rho^{-(0)} e^+ \nu_e$				
$r_V$	5.0% <sub>stat.</sub> 4.0% <sub>syst.</sub>	2.0% <sub>stat.</sub> 2.0% <sub>syst.</sub>		
$r_A$	8.0% <sub>stat.</sub> 4.0% <sub>syst.</sub>	3.0% <sub>stat.</sub> 2.0% <sub>syst.</sub>		

STCF will largely improve the precisions of the form factors over all the modes

# Form factors $f_+^{D \rightarrow h}$



Crucial tests on the LQCD results

# Lepton Flavor Universality



LFU is **critical** to test the SM and search for new physics beyond the SM

Pure-leptonic modes

$$R_{D(s)^+} = \frac{\Gamma(D(s)^+ \rightarrow \tau^+ \nu_\tau)}{\Gamma(D(s)^+ \rightarrow \mu^+ \nu_\mu)} = \frac{m_{\tau^+}^2 \left(1 - \frac{m_{\tau^+}^2}{m_{D(s)^+}^2}\right)^2}{m_{\mu^+}^2 \left(1 - \frac{m_{\mu^+}^2}{m_{D(s)^+}^2}\right)^2}$$

Semi-leptonic modes

$$R_{\mu/e} = \Gamma_{D^0 \rightarrow K^- \mu^+ \nu_\mu} / \Gamma_{D^0 \rightarrow K^- e^+ \nu_e}$$



1 $\sigma$  difference?

2 $\sigma$  difference?


2.93/fb@3773MeV;  
3.19/fb @ 4178MeV

	$R(D_s^+)$	$R(D^+)$	$R(K^-)$	$R(\bar{K}^0)$	$R(\pi^-)$	$R(\pi^0)$
SM	9.74(1)	2.66(1)	0.975(1)	0.975(1)	0.985(2)	0.985(2)
BESIII	10.19(52)	3.21(64)	0.974(14)	1.013(29)	0.922(37)	0.964(45)

- Large uncertainty from BESIII, dominant by statistically limited
- BESIII and STCF would improve them significantly

# Determination of $\gamma/\phi_3$ angle


- The **cleanest way** to extract  $\gamma$  is from  **$B \rightarrow DK$**  decay



- Interference between tree-level decays; theoretically clean
- current uncertainty  $\sigma(\gamma) \sim 5^\circ$
- however, theoretical relative error  $\sim 10^{-7}$  (very small!)

- Information of  **$D$  decay strong phase** is needed

- Best way is to employ **quantum coherence** **DD production** at threshold



Prog. Theor. Exp. Phys. 2020 083C01 (2020)

# Determination of $\gamma/\phi_3$ angle

Runs	Collected / Expected integrated luminosity	Year attained	$\gamma/\phi_3$ sensitivity
LHCb Run-1 [7, 8 TeV]	3 fb <sup>-1</sup>	2012	8°
LHCb Run-2 [13 TeV]	5 fb <sup>-1</sup>	2018	4°
Belle II Run	50 ab <sup>-1</sup>	2025	1.5°
LHCb upgrade I [14 TeV]	50 fb <sup>-1</sup>	2030	< 1°
LHCb upgrade II [14 TeV]	300 fb <sup>-1</sup>	(>)2035	< 0.4°

BESIII 20/fb:

$\sigma(\gamma) \sim 0.4^\circ$


STCF is needed!

Three methods for exploiting interference  
(choice of  $D^0$  decay modes):

- ❑ GLW: Use CP eigenstates of  $D^{(*)0}$  decay, e.g.  $D^0 \rightarrow K_S \pi^0$ ,  $D^0 \rightarrow \pi^+ \pi^-$
- ❑ ADS: Use doubly Cabibbo-suppressed decays, e.g.  $D^0 \rightarrow K^+ \pi^-$ 
  - With 1 ab<sup>-1</sup> @ STCF :  $\sigma(\cos\delta_{K\pi}) \sim 0.007$ ;  $\sigma(\delta_{K\pi}) \sim 2^\circ \rightarrow \sigma(\gamma) < 0.5^\circ$
- ❑ BPGGSZ: Use Dalitz plot analysis of 3-body  $D^0$  decays, e.g.  $K_S \pi^+ \pi^-$ ; high statistics; need precise Dalitz model
  - STCF reduces the contribution of  $D$  Dalitz model to a level of  $\sim 0.1^\circ$

Decay mode	Quantity of interest
$D \rightarrow K_S^0 \pi^+ \pi^-$	$c_i$ and $s_i$
$D \rightarrow K_S^0 K^+ K^-$	$c_i$ and $s_i$
$D \rightarrow K^\pm \pi^\mp \pi^+ \pi^-$	$R, \delta$
$D \rightarrow K^+ K^- \pi^+ \pi^-$	$c_i$ and $s_i$
$D \rightarrow \pi^+ \pi^- \pi^+ \pi^-$	$F_+$ or $c_i$ and $s_i$
$D \rightarrow K^\pm \pi^\mp \pi^0$	$R, \delta$
$D \rightarrow K_S^0 K^\pm \pi^\mp$	$R, \delta$
$D \rightarrow \pi^+ \pi^- \pi^0$	$F_+$
$D \rightarrow K_S^0 \pi^+ \pi^- \pi^0$	$F_+, c_i$ and $s_i$
$D \rightarrow K^+ K^- \pi^0$	$F_+$
$D \rightarrow K^\pm \pi^\mp$	$\delta$

# Determination of $\gamma/\phi_3$ angle in CKM



STCF will provide complementary information on the strong phase and allow detailed comparisons in different models



# $D^0$ - $\bar{D}^0$ mixing and CPV at STCF

STCF provide **a unique place** for the study of  $D^0$ - $\bar{D}^0$  mixing and CPV by means of **quantum coherence** of  $D^0$  and  $\bar{D}^0$  produced through

$$\psi(3770) \rightarrow (D^0 \bar{D}^0)_{\text{CP}=-} \text{ or } \psi(4140) \rightarrow D^0 \bar{D}^{*0} \rightarrow \pi^0 (D^0 \bar{D}^0)_{\text{CP}=-} \text{ or } \gamma (D^0 \bar{D}^0)_{\text{CP}=+}$$

- Mixing rate  $R_M = \frac{x^2+y^2}{2} \sim \mathbf{10^{-5}}$  with 1  $\text{ab}^{-1}$  data at 3.773 GeV via **same charged** final states  $(K^\pm \pi^\mp)(K^\pm \pi^\mp)$  or  $(K^\pm l^\mp \nu)(K^\pm l^\mp \nu)$
- Mixing parameters and CPV parameters with 1  $\text{ab}^{-1}$  data at 4009 MeV via coherent (C-even and C-odd) and incoherent process
- $\Delta A_{CP} \sim \mathbf{10^{-3}}$  for KK and  $\pi\pi$  channels


# Precision estimation at STCF

	1/ab @4009 MEV (only QC   QC+incoherent) (preliminary estimation)		BELLEII(50/ab) [PTEP2019, 123C01]	LHCb(50/fb) (SL   Prompt) [arXiv:1808.08865]	
$x(\%)$	0.036	0.035	0.03	0.024	0.012
$y(\%)$	0.023	0.023	0.02	0.019	0.013
$r_{CP}$	0.017	0.013	0.022	0.024	0.011
$\alpha_{CP}(^\circ)$	1.3	1.0	1.5	1.7	0.48

- The only QC results: contains  $D^0 \rightarrow K_S \pi \pi$ ,  $D^0 \rightarrow K^- \pi^+ \pi^0$  and general CP tag decay channels
- The QC+incoherent results: combines coherent and incoherent  $D^0$  meson samples
- The BELLE II and LHCb results only contain incoherent  $D^0 \rightarrow K_S \pi \pi$  channel



# Charmed baryon decays




	Structure	$J^P$	Mass, MeV	Width, MeV	Decay
$\Lambda_c^+$	$udc$	$(1/2)^+$	$2286.46 \pm 0.14$	$(200 \pm 6)$ fs	weak
$\Xi_c^+$	$usc$	$(1/2)^+$	$2467.8^{+0.4}_{-0.6}$	$(442 \pm 26)$ fs	weak
$\Xi_c^0$	$dsc$	$(1/2)^+$	$2470.88^{+0.34}_{-0.8}$	$112^{+13}_{-10}$ fs	weak
$\Sigma_c^{++}$	$uuc$	$(1/2)^+$	$2454.02 \pm 0.18$	$2.23 \pm 0.30$	$\Lambda_c^+ \pi^+$
$\Sigma_c^+$	$udc$	$(1/2)^+$	$2452.9 \pm 0.4$	$< 4.6$	$\Lambda_c^+ \pi^0$
$\Sigma_c^0$	$ddc$	$(1/2)^+$	$2453.76 \pm 0.18$	$2.2 \pm 0.4$	$\Lambda_c^+ \pi^-$
$\Xi_c'^+$	$usc$	$(1/2)^+$	$2575.6 \pm 3.1$	—	$\Xi_c^+ \gamma$
$\Xi_c'^0$	$dsc$	$(1/2)^+$	$2577.9 \pm 2.9$	—	$\Xi_c^0 \gamma$
$\Omega_c^0$	$ssc$	$(1/2)^+$	$2695.2 \pm 1.7$	$(69 \pm 12)$ fs	weak
$\Sigma_c^{*++}$	$uuc$	$(3/2)^+$	$2518.4 \pm 0.6$	$14.9 \pm 1.9$	$\Lambda_c^+ \pi^+$
$\Sigma_c^{*+}$	$udc$	$(3/2)^+$	$2517.5 \pm 2.3$	$< 17$	$\Lambda_c^+ \pi^0$
$\Sigma_c^{*0}$	$ddc$	$(3/2)^+$	$2518.0 \pm 0.5$	$16.1 \pm 2.1$	$\Lambda_c^+ \pi^-$
$\Xi_c^{*+}$	$usc$	$(3/2)^+$	$2645.9^{+0.5}_{-0.6}$	$< 3.1$	$\Xi_c \pi$
$\Xi_c^{*0}$	$dsc$	$(3/2)^+$	$2645.9 \pm 0.5$	$< 5.5$	$\Xi_c \pi$
$\Omega_c^{*0}$	$ssc$	$(3/2)^+$	$2765.9 \pm 2.0$	—	$\Omega_c^0 \gamma$

# Prospects at BESIII

4.6 - 4.9 GeV	Charmed baryon/ $XYZ$ cross-sections	$0.56 \text{ fb}^{-1}$ at 4.6 GeV	$15 \text{ fb}^{-1}$ at different $\sqrt{s}$
4.74 GeV	$\Sigma_c^+ \Lambda_c^-$ cross-section	N/A	$1.0 \text{ fb}^{-1}$
4.91 GeV	$\Sigma_c \bar{\Sigma}_c$ cross-section	N/A	$1.0 \text{ fb}^{-1}$
4.95 GeV	$\Xi_c$ decays	N/A	$1.0 \text{ fb}^{-1}$

	Leading hadronic decay	Typical two-body decay	Leading SL decay
$\Lambda_c^+$	$\mathcal{B}(K^- p \pi^+) =$ 2014: $(5.0 \pm 1.3)\%$ (26%) 2017(w/ BESIII): $(6.35 \pm 0.33)\%$ (5.2%) 5 $\text{fb}^{-1}$ : $\frac{\delta \mathcal{B}}{\mathcal{B}} < 2\%$	$\mathcal{B}(K_S^0 p) =$ 2014: $(1.2 \pm 0.3)\%$ (26%) BESIII: $(1.52 \pm 0.08)\%$ (5.6%) 5 $\text{fb}^{-1}$ : $\frac{\delta \mathcal{B}}{\mathcal{B}} < 2\%$	$\mathcal{B}(\Lambda e^+ \nu_e) =$ 2014: $(2.1 \pm 0.6)\%$ (29%) BESIII: $(3.63 \pm 0.43)\%$ (12%) 5 $\text{fb}^{-1}$ : $\frac{\delta \mathcal{B}}{\mathcal{B}} \sim 3.3\%$
$D^0$	$\mathcal{B}(K^- \pi^+) = (3.89 \pm 0.04)\%$ (1.0%)	$\mathcal{B}(K_S^0 \pi^0) = (1.19 \pm 0.04)\%$ (3.4%)	$\mathcal{B}(K^- e^+ \nu_e) = (3.53 \pm 0.03)\%$ (0.8%)
$D^+$	$\mathcal{B}(K^- \pi^+ \pi^+) = (8.98 \pm 0.28)\%$ (3.1%)	$\mathcal{B}(K_S^0 \pi^+) = (1.47 \pm 0.08)\%$ (5.4%)	$\mathcal{B}(K_S^0 e^+ \nu_e) = (4.41 \pm 0.07)\%$ (1.5%)
$D_s^+$	$\mathcal{B}(K^- K^+ \pi^+) = (5.45 \pm 0.17)\%$ (3.8%)	$\mathcal{B}(K_S^0 K^+) = (1.40 \pm 0.05)\%$ (3.6%)	$\mathcal{B}(\phi e^+ \nu_e) = (2.39 \pm 0.23)\%$ (9.6%)



Mode	Expected rate (%)	Relative uncertainty (%)
$\Lambda_c^+ \rightarrow \Lambda l^+ \nu$	3.6 [94, 95]	3.3
$\Lambda_c^+ \rightarrow \Lambda^* l^+ \nu$	0.7 [96, 97]	10
$\Lambda_c^+ \rightarrow N K e^+ \nu_e$	0.7 [96]	10
$\Lambda_c^+ \rightarrow \Sigma \pi l^+ \nu$	0.7 [96]	10
$\Lambda_c^+ \rightarrow n e^+ \nu_e$	0.2 [94, 98, 99]	17

first measurement

# STCF: precision studies of the charmed baryon decays

Era of precision study of the charmed baryon ( $\Lambda_c$ ,  $\Xi_c$  and  $\Omega_c$ ) decays  
to help developing more reliable QCD-derived models in charm sector

- ▣ **Hadronic decays:**

to explore as-yet-unmeasured channels and understand full picture of intermediate structures in charmed baryon decays, esp., those with neutron/ $\Sigma$ / $\Xi$  particles

- ▣ **Semi-leptonic decays:**

to test LQCD calculations and LFU

- ▣ **CPV in charmed baryon:** BP and BV two-body decay asymmetry, charge-dependent rate of SCS

- ▣ **Charmed Baryons Spectroscopy** : (63 P-wave states from QM, less than 20 are observed!)

- ▣ **Rare decays:** LFV, BNV, FCNC

*STCF will provide very precise measurements of their overall decays, up to the unprecedented level of  $10^{-6} \sim 10^{-7}$*

# Summary

- **BESIII and STCF** are the crucial **precision frontier**
- Important playground for studying non-perturbative **QCD, constrain EW theory** and test the SM
  - ✓ CKM matrix elements
  - ✓ LQCD calibration
  - ✓ Search for violation of LFU in charm sector
  - ✓ Precision measurement of the charmed baryon decays
- Complementary to Belle II and LHCb in understanding the QCD/EW models and searching for new physics



Thank you!

谢谢!

# Data samples

Data samples with  $1 \text{ ab}^{-1}$  integral luminosity

Data Set	STCF					Belle II		
	process	$\sigma/\text{nb}$	N	ST eff./%	ST N	$\sigma/\text{nb}$	N	Tag N
$J/\psi$	—	—	$1.0 \times 10^{12}$	—	—	—	—	—
$\psi(2S)$	—	—	$3.0 \times 10^{11}$	—	—	—	—	—
$D^0$	$D^0 \bar{D}^0(3.77)$	$\sim 3.6$	$3.6 \times 10^9$	10.8	$0.78 \times 10^9$	—	$1.4 \times 10^9$	—
$D^+$	$D^+ D^-(3.77)$	$\sim 2.8$	$2.8 \times 10^9$	9.4	$0.53 \times 10^9$	—	$7.7 \times 10^8$	—
$D_s$	$D_s D_s^*(4.18)$	$\sim 0.9$	$0.9 \times 10^9$	6.0	$0.11 \times 10^9$	—	$2.5 \times 10^8$	—
$\tau^+$	$\tau^+ \tau^-(3.68)$	$\sim 2.4$	$2.4 \times 10^9$	—	—	0.9	$0.9 \times 10^9$	—
	$\tau^+ \tau^-(4.25)$	$\sim 3.6$	$3.5 \times 10^9$	—	—	—	—	—
$\Lambda_c$	$\Lambda_c \Lambda_c(4.64)$	$\sim 0.6$	$5.5 \times 10^8$	5.0	$0.55 \times 10^8$	—	$1.6 \times 10^8$	$3.6 \times 10^{4*}$

The luminosity is  $1.0 \text{ ab}^{-1}$ . \* process  $e^+ e^- \rightarrow D^{(*)-} \bar{p} \pi^+ \Lambda_c^+$ .

- Belle-II (50/ab) has 50~100 times more statistics
- STCF is expected to have higher detection efficiency
- STCF has low backgrounds for productions at threshold

# $D^0$ mixing and CPV parameters

- Three kinds of  $D^0\bar{D}^0$  samples can be used @4009MeV
  - **Quantum-incoherent flavor specific  $D^0$  samples:**  $D^{*+} \rightarrow \textcolor{red}{D}^0\pi^+$ 
    - Help to improve precision of strong-phase difference measurement
    - Be used to constrain the charm mixing and CPV parameters
  - **Quantum-coherent C-even  $D^0\bar{D}^0$  samples:**  $D^{*0}\bar{D}^0 \rightarrow \textcolor{red}{D}^0\bar{\textcolor{red}{D}}^0\gamma$ 
    - Be used to perform charm mixing and CPV parameters measurements
      - The interference effect, containing mixing and CPV, is doubled compare to incoherent case
    - Help to constrain the strong-phase difference and CP fraction measurements
  - **Quantum-coherent C-odd  $D^0\bar{D}^0$  samples:**  $D^{*0}\bar{D}^0 \rightarrow \textcolor{red}{D}^0\bar{\textcolor{red}{D}}^0\pi^0$ 
    - Same as  $D^0\bar{D}^0$  samples @3770, improve precision of strong-phase difference measurements and CP fraction measurements

	BES III $10^{33} \text{ cm}^{-2} \text{ s}^{-1}$	STCF $10^{35} \text{ cm}^{-2} \text{ s}^{-1} (1 \text{ ab}^{-1})$
$J/\psi$	$10 \times 10^9$	$10 \times 10^{11}$
$\psi(2S)$	$3 \times 10^9$	$3 \times 10^{11}$
$D(3.773 \text{ GeV})$	$6 \times 10^7$	$6 \times 10^9$
$D_s (4.17 \text{ GeV})$	$1 \times 10^7$	$1 \times 10^9$
$\tau^+ \tau^- (3.68 \text{ GeV})$		$2.4 \times 10^9$
$\tau^+ \tau^- (4.25 \text{ GeV})$	$3 \times 10^7$	$3.5 \times 10^9$
$\Lambda_c^+ \Lambda_c^- (4.64 \text{ GeV})$	$3 \times 10^6$	$6 \times 10^8$



# Charm mixing and CPV parameters

- For quantum-incoherent  $D^0$  meson samples[1]:

$$\overline{P}'(m_{12}^2, m_{13}^2) = a_0 \overline{P} + a_1 r_{c\overline{P}}^{-2} P + r_{c\overline{P}}^{-1} \sqrt{P\overline{P}} (C^- a_2 + S^- a_3)$$

- For quantum-coherent  $D^0 \overline{D}^0$  samples:

$$\begin{aligned} r_{c\overline{P}} e^{i\alpha_{c\overline{P}}} &= \frac{q}{p}, \\ S^\pm &= \sin(\Delta\delta_D \pm \alpha_{c\overline{P}}), \\ C^\pm &= \cos(\Delta\delta_D \pm \alpha_{c\overline{P}}), \\ a_0 &= \frac{1}{2} \left( \frac{1}{1-y_D^2} + \frac{1}{1+x_D^2} \right) \\ &= 1 + \frac{1}{2} (-x_D^2 + y_D^2) + O((x_D + y_D)^3), \\ a_1 &= \frac{1}{2} \left( \frac{1}{1-y_D^2} - \frac{1}{1+x_D^2} \right) \\ &= \frac{1}{2} (x_D^2 + y_D^2) + O((x_D + y_D)^3), \\ a_2 &= \frac{y_D}{1-y_D^2} = y_D + O((x_D + y_D)^3), \\ a_3 &= \frac{x_D}{1+x_D^2} = x_D + O((x_D + y_D)^3). \end{aligned}$$

$$P_{corr}^C((m_{12}^2)_1, (m_{13}^2)_1, (m_{12}^2)_2, (m_{13}^2)_2) = b_0^C \left[ P_1 \overline{P}_2 + \overline{P}_1 P_2 + 2C \sqrt{P_1 \overline{P}_1 P_2 \overline{P}_2} (C_1 C_2 + S_1 S_2) \right]$$

$$\begin{aligned} &+ b_1^C \left[ r_{c\overline{P}}^{-2} P_1 P_2 + r_{c\overline{P}}^2 \overline{P}_1 \overline{P}_2 + 2C \sqrt{P_1 \overline{P}_1 P_2 \overline{P}_2} (C_1^+ C_2^+ - S_1^+ S_2^+) \right] \\ &+ b_2^C \left[ \sqrt{P_2 \overline{P}_2} C_2^+ (r_{c\overline{P}} \overline{P}_1 + r_{c\overline{P}}^{-1} P_1) + C \sqrt{P_1 \overline{P}_1} C_1^+ (r_{c\overline{P}} \overline{P}_2 + r_{c\overline{P}}^{-1} P_2) \right] \\ &+ b_3^C \left[ \sqrt{P_2 \overline{P}_2} S_2^+ (r_{c\overline{P}} \overline{P}_1 - r_{c\overline{P}}^{-1} P_1) + C \sqrt{P_1 \overline{P}_1} S_1^+ (r_{c\overline{P}} \overline{P}_2 - r_{c\overline{P}}^{-1} P_2) \right], \end{aligned}$$

$$\begin{aligned} b_0^C &= \frac{1}{2} \left[ \frac{1 + C y_D^2}{(1 - y_D^2)^2} + \frac{1 - C x_D^2}{(1 + x_D^2)^2} \right] \approx a_0 + \frac{C+1}{2} (-x_D^2 + y_D^2) \\ b_1^C &= \frac{1}{2} \left[ \frac{1 + C y_D^2}{(1 - y_D^2)^2} - \frac{1 - C x_D^2}{(1 + x_D^2)^2} \right] \approx (C+2) a_1, \\ b_2^C &= \frac{(1+C) y_D}{(1 - y_D^2)^2} \approx (1+C) a_2, \\ b_3^C &= \frac{(1+C) x_D}{(1 + x_D^2)^2} \approx (1+C) a_3, \end{aligned}$$


# Integral Luminosity of STCF

- No Synchrotron radiation mode, assume running time 9 months/year
- Assume data taking efficiency 90%

$$0.5 \times 10^{35} \text{cm}^{-2} \text{s}^{-1} \times 86400 \text{s} \times 270 \text{days} \times 90\% \sim \mathbf{1.0 ab^{-1} / year}$$

10 years data taking, total 20  $\text{ab}^{-1}$  conservatively

Excellent opportunities for the  $\tau$ -charm physics





## at Belle II

► each 1  $\text{ab}^{-1}$  dataset provides

- $\sim 1.1 \times 10^9 B\bar{B} \Rightarrow$  a **B-factory**;
- $\sim 1.3 \times 10^9 c\bar{c} \Rightarrow$  a **charm factory**;
- $\sim 0.9 \times 10^9 \tau^+\tau^- \Rightarrow$  a  **$\tau$  factory**;
- wide  $E_{\text{CM}}^{\text{eff}} = [0.5-10] \text{ GeV}$  via ISR.

# Features in studying charm hadron decays

	BESIII/ STCF	Belle II	LHCb
Production yields	★★	★★★★★	★★★★★★
Background level	★★★★★★	★★★★	★★
Systematic error	★★★★★★	★★★★	★★
Completeness	★★★★★★	★★★★	★
(Semi)-Leptonic mode	★★★★★★	★★★★★	★★
Neutron/ $K_L$ mode	★★★★★★	★★★★	☆
Photon-involved	★★★★★★	★★★★★	★
Absolute measurement	★★★★★★	★★★★	☆



- Most are **precision** measurements, which are mostly dominant by the **systematic** uncertainty
- STCF has **overall advantages** in several studies